

SHORT GENOME REPORT

Open Access

Draft genome sequence of *Bacillus azotoformans* MEV2011, a (Co-) denitrifying strain unable to grow with oxygen

Maja Nielsen^{1†}, Lars Schreiber^{2†}, Kai Finster^{1,3} and Andreas Schramm^{1,2*}

Abstract

Bacillus azotoformans MEV2011, isolated from soil, is a microaerotolerant obligate denitrifier, which can also produce N₂ by co-denitrification. Oxygen is consumed but not growth-supportive. The draft genome has a size of 4.7 Mb and contains key genes for both denitrification and dissimilatory nitrate reduction to ammonium.

Keywords: *Bacillus azotoformans*, Denitrification, Codenitrification, Oxygen

Introduction

Species of the genus *Bacillus* are characterized as Gram-positive, facultative aerobic bacteria capable of forming endospores [1]. In the absence of oxygen, many *Bacillus* species can respire with nitrate instead, employing either dissimilatory nitrate reduction to ammonium or denitrification [2,3]. Despite the widespread occurrence of nitrate-reducing bacilli, their molecular and genetic basis remained poorly investigated [4,5]. Only recently, genome sequencing of two denitrifying type strains, *B. azotoformans* LMG 9581^T and *B. bataviensis* LMG 21883^T, has yielded first insights into the genomic inventory of nitrate reduction and denitrification in Gram-positives [6].

Classification and features

B. azotoformans MEV2011 (Figure 1) was isolated at 28°C on anoxic King B plates [7] amended with KNO₃ (5 g L⁻¹) from a highly diluted top soil sample at Aarhus University, Denmark. Strain MEV2011 resembles the type strain in its chemoorganotrophic growth on short-chain fatty acids, complete denitrification, and absence of fermentation [8]. However, it differs from the type strain by its inability to grow with oxygen, even though it can tolerate and consume oxygen at atmospheric concentrations. Growth by denitrification (verified by

¹⁵N incubations; data not shown) starts at microaerobic conditions (<30 μM O₂; Figure 2), yet the initial presence of oxygen in the growth medium leads to longer lag phases and no increase in final density of the culture (Figure 3); growth without nitrate was never observed. Therefore, we characterize *B. azotoformans* MEV2011 as microaerotolerant obligate denitrifier. In addition, *B. azotoformans* MEV2011 is capable of co-denitrification, a co-metabolic process, in which reduced nitrogen compounds like amino acids or hydroxylamine react with NO⁺ formed during denitrification to produce N₂O or N₂ [9]; co-denitrification was verified by the mass spectrometric detection of ³⁰N₂ + ²⁹N₂ in cultures growing on tryptic soy broth (TSB) and ¹⁵NO₃⁻, as suggested in [9]. *B. azotoformans* MEV2011 is available from the BCCM/LMG Bacteria Collection as strain LMG 28302; its general features are summarized in Table 1.

Genome sequencing and annotation

Genome project history

Bacillus azotoformans MEV2011 was selected for whole genome sequencing based on its unusual “obligate” denitrifying phenotype, i.e. its inability to grow under oxic conditions, together with its co-denitrifying capacity. Comparing the genome of strain MEV2011 to that of the oxygen-respiring and conventionally denitrifying type strain [8] may provide insights into the molecular basis of its metabolic features. The draft genome

* Correspondence: andreas.schramm@biology.au.dk

†Equal contributors

¹Section for Microbiology, Department of Bioscience, Aarhus University, Aarhus, Denmark

²Center for Geomicrobiology, Department of Bioscience, Aarhus University, Aarhus, Denmark

Full list of author information is available at the end of the article




Figure 1 Phylogenetic tree highlighting the position of *Bacillus azotoformans* MEV2011 (shown in red) relative to closely related ($\geq 95\%$ sequence similarity) type strains within the *Bacillaceae*. Pre-aligned sequences were retrieved from the Ribosomal Database Project (RDP) [37]. Alignment of the *B. azotoformans* MEV2011 sequence as well as manual alignment optimization was performed in ARB [38]. The maximum likelihood tree was inferred from 1,478 aligned positions of 16S rRNA gene sequences and calculated based on the General Time Reversible (GTR) model with gamma rate heterogeneity using RAxML 7.4.2 [39]. Type strains with corresponding published genomes are shown in bold face. Open and closed circles indicate nodes with bootstrap support (1,000 replicates) of 50-80% and $>80\%$, respectively. *Escherichia coli* ATCC 11577^T (X80725) was used to root the tree (not shown). Scale bar, 0.1 substitutions per nucleotide position.

sequence was completed on July 20, 2013. The genome project is deposited in the Genomes OnLine Database (GOLD) as project Gp0043190. Raw sequencing reads have been deposited at the NCBI Sequence Read Archive (SRA) under the experiment numbers SRX527325 (100 bp library) and SRX527326 (400 bp

library). This Whole Genome Shotgun project has been deposited at GenBank under the accession number JJRY00000000. The version described in this paper is version 1. Table 2 presents the project information and its association with MIGS version 2.0 compliance [27].




Figure 2 Consumption of oxygen (\square ; measured online with an oxygen microsensor) and nitrate (\bullet ; measured by HPLC) during growth (\blacktriangle ; OD_{600}) of *B. azotoformans* MEV2011. No growth was observed at oxygen concentrations >30 - $35 \mu\text{M}$, and the initiation of growth coincided with the first detection of $^{30}\text{N}_2$ from $^{15}\text{NO}_3^-$ (data not shown), indicating that growth was coupled to denitrification.




Figure 3 Length of lag phase (h; bars), and final biomass (OD_{600} ; circles) of *B. azotoformans* MEV2011 as function of the initial oxygen concentration in the culture. Cultures were grown in TSB (10 g L^{-1} , Scharlau®) amended with 3 mM KNO_3 . Black and grey bars and circles represent data from replicate incubations. Growth was first detected when oxygen had been consumed to $<30\text{--}35 \mu\text{M}$ (see Figure 2), explaining the increasing lag time with increasing oxygen concentrations. The final OD was almost identical in all incubations and unrelated to the initial oxygen concentration, indicating that oxygen did not contribute to biomass production.

Growth conditions and genomic DNA preparation

B. azotoformans MEV2011 was grown at 28°C in N_2 -flushed TSB (10 g L^{-1} , Scharlau®) amended with KNO_3 (3 g L^{-1}). DNA was extracted using the DNeasy Blood & Tissue kit (Qiagen®).

Genome sequencing and assembly

Sequencing of the *B. azotoformans* MEV2011 genome was performed with an Ion Torrent PGM sequencer (Life Sciences) using 100 and 400 bp sequencing chemistries. Sequencing libraries were prepared using Ion Xpress™ Plus Fragment Library Kits (Life Sciences), and Ion OneTouch™ Template Kits (Life Sciences). Sequencing of the 100 bp library generated 442,853 reads (representing 42 Mbp of sequence information), while sequencing of the 400 bp library generated 2,401,947 reads (477 Mbp). Together, both libraries achieved a genome coverage of $c. 110\times$ for an estimated genome size of 4.7 Mbp. The reads were quality trimmed using the prinseq-lite.pl script [11] with the following parameters; reads generated with 100 bp chemistry: `-min_len 50 -trim_to_len 110 -trim_left 15 -trim_qual_right 20 -trim_qual_window 4 -trim_qual_type mean`; reads generated with 400 bp chemistry: `-min_len 50 -trim_to_len 400 -trim_left 15 -trim_qual_right 20 -trim_qual_window 4 -trim_qual_type mean`. The trimmed reads (2,491,456 reads representing 444 Mbp) were assembled using MIRA 3.9.18 [12] with the following parameters: `job = genome,denovo,accurate; technology = iontor`. In parallel, the reads were also assembled using Newbler 2.6 (Roche) with the following parameters: `-mi 96 -ml 50` (i.e. 96% minimum sequence similarity and 50 bp minimum overlap). Contigs shorter than 1,000 bp were removed from both assemblies. All remaining

contigs were trimmed by 50 bp from the 5' and the 3' ends using the prinseq-lite.pl script in order to remove error-prone contig ends. The two assemblies were merged and manually inspected using Sequencher 5.0.1 (Genecodes). In cases where the bases of the two assemblies disagreed, the Newbler variant was preferred. Contigs not contained in both assemblies were removed from the data set. The final assembly yielded 56 contigs representing 4.7 Mbp of sequence information.

Genome annotation

The draft genome was auto-annotated using the standard operation procedure of the Integrated Microbial Genomes Expert Review (IMG-ER) platform developed by the Joint Genome Institute, Walnut Creek, CA, USA [13]. In short, CRISPR regions were identified by CRT [14] and PILERCR [15], tRNAs were identified by tRNAScan-SE-1.23 [16], rRNAs were identified by RNAmmer [17], and finally all other genes were identified by Prodigal [18]. Functional annotation was based on gene comparisons with the KEGG database (release 63.0, July 1, 2012) [19], the PFAM database (version 25.0, March 30, 2011) [20], the cluster of orthologous groups (COG) [21] database, and the TIGRFam database (release 11.0, August 3, 2011) [22].

Genome properties

The MEV2011 draft genome is 4,703,886 bp long and comprises 56 contigs ranging in size from 1,773 to 525,568 bp, with an overall GC content of 37.49% (Table 3). Of the 4,986 predicted genes, 4,809 (96.45%) are protein-coding genes, and 177 are RNAs. Of the RNAs, 94 are tRNAs, and 37 are rRNAs. The number of 5S rRNAs as well as the number of partial 16S and 23S

Table 1 Classification and general features of *Bacillus azotoformans* MEV2011 [27]

MIGS ID	Property	Term	Evidence code ^a
	Classification	Domain <i>Bacteria</i>	TAS [28]
		Phylum <i>Firmicutes</i>	TAS [29–31]
		Class <i>Bacilli</i>	TAS [32,33]
		Order <i>Bacillales</i>	TAS [29,34]
		Family <i>Bacillaceae</i>	TAS [29,35]
		Genus <i>Bacillus</i>	TAS [29,36]
		Species <i>Bacillus azotoformans</i>	TAS [8]
		Strain: MEV2011 (LMG 28302)	IDA
	Gram stain	Variable	IDA
	Cell shape	Rod	IDA
	Motility	Motile	IDA
	Sporulation	Endospore-forming	IDA
	Temperature range	15 – 42°C	IDA
	Optimum temperature	39 – 42°C	IDA
	pH range; Optimum	4–9; 7	IDA
	Carbon source	Malate, acetate, lactate, citrate, succinate, yeast extract	IDA
	Terminal electron acceptor	Nitrate, nitrite, NO, N ₂ O (O ₂ is reduced but does not support growth)	IDA
MIGS-6	Habitat	Soil	IDA
MIGS-6.3	Salinity	0–3% NaCl (w/v)	IDA
MIGS-22	Oxygen requirement	Anaerobic, microaerotolerant	IDA
MIGS-15	Biotic relationship	Free-living	IDA
MIGS-14	Pathogenicity	Non-pathogen	IDA
MIGS-4	Geographic location	Denmark/Aarhus University campus, Aarhus	IDA
MIGS-5	Sample collection	2011-02-01	IDA
MIGS-4.1	Latitude	56° 10' 0.12" N	IDA
MIGS-4.2	Longitude	10° 12' 6.12" E	IDA
MIGS-4.4	Altitude	38.6 m	IDA

^aEvidence codes - IDA: Inferred from Direct Assay; TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [10].

rRNA genes indicates a total of 11 rRNA operons. Most (75.3%) protein-coding genes were assigned to putative functions. The distribution of genes into COG functional categories is presented in Table 4.

Insights from the genome sequence

Overall, the genome of the novel strain MEV2011 appeared highly similar to that of the *B. azotoformans* type strain LMG 9581^T [8]. *In silico* DNA–DNA hybridization (DDH) was performed for the assembled MEV2011 genome against the published genome of LMG 9581^T (Acc. number NZ_AJLR00000000); the contigs of *B. azotoformans* LMG 9581^T were assembled into one FASTA file before uploading to the online genome-to-genome calculator provided by the DSMZ [23]. Using the GGDC 2.0 model, DHH estimates were always >70%, irrespective of the formula used for computing DHH,

and with probabilities between 78 and 87%. These results confirm that MEV2011 is a novel strain of the species *B. azotoformans*.

Just as *B. azotoformans* LMG 9581^T, strain MEV2011 carries multiple copies of key denitrification genes, encodes both membrane-bound and periplasmic nitrate reductases, and the key genes for nitrite reduction to both NO (in denitrification) and ammonium (in DNRA); see (Additional file 1: Table S1) and reference [6] for details. Modularity and redundancy in nitrate reduction pathways has also been observed in other *Bacillus* species (e.g. *B. bataviensis* [6], *Bacillus* sp. strain ZYK [24], *Bacillus* sp. strain 1NLA3E [25]), and may be a general feature of nitrate-reducing members of this genus.

All genes essential for aerobic respiration were identified, including those for terminal oxidases (see Additional

Table 2 Project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	High quality draft
MIGS-28	Libraries used	IonTorrent 100 bp and 400 bp single end reads
MIGS-29	Sequencing platforms	IonTorrent PGM
MIGS-31.2	Fold coverage	110x
MIGS-30	Assemblers	Newbler 2.6, MIRA 3.9.18, Sequencher 5.0.1
MIGS-32	Gene calling method	Prodigal
	Locus Tag	M670
	Genbank ID	JJRY00000000
	Genbank Date of Release	2014-06-16
	GOLD ID	Gi0050495
	BIOPROJECT	PRJNA209301
	Project relevance	Environmental, co-denitrification
MIGS 13	Source Material Identifier	LMG 28302

file 1: Table S2) and for detoxifying reactive oxygen species (see Additional file 1: Table S3). Therefore, the inability of *B. azotoformans* MEV2011 to grow with oxygen remains a conundrum and in some way resembles that of various sulfate-reducing bacteria, which also consume oxygen and even produce ATP during oxic respiration but are unable to grow in the presence of oxygen [26].

Conclusion

Based on our whole genome comparison, the microaerophilic obligate (co-) denitrifying *Bacillus* sp. MEV2011

Table 3 Nucleotide content and gene count levels of the genome

Attribute	Value	% of total ^a
Genome size (bp)	4,703,886	100
DNA coding (bp)	4,075,859	86.7
DNA G + C (bp)	1,763,498	37.5
DNA scaffolds	56	100
Total genes	4,986	100
Protein coding genes	4,809	96.5
RNA genes	177	3.6
Pseudo genes	0	0
Genes in internal clusters	3,448	69.1
Genes with function prediction	3,755	75.3
Genes assigned to COGs	2,809	56.3
Genes with Pfam domains	3,890	78.0
Genes with signal peptides	182	3.7
Genes with transmembrane helices	1,233	24.7
CRISPR repeats	4	-

^aThe total is based on either the size of the genome in base pairs or the total number of protein coding genes in the annotated genome.

Table 4 Number of genes associated with general COG functional categories

Code	Value	% age ^a	Description
A	0	0	RNA processing and modification
J	160	3.32	Translation, ribosomal structure and biogenesis
K	246	5.11	Transcription
L	185	3.85	Replication, recombination and repair
B	1	0.02	Chromatin structure and dynamics
D	34	0.70	Cell cycle control, Cell division, chromosome partitioning
V	45	0.94	Defense mechanisms
T	207	4.30	Signal transduction mechanisms
M	123	2.56	Cell wall/membrane biogenesis
N	77	1.60	Cell motility
U	50	1.04	Intracellular trafficking and secretion
O	106	2.20	Posttranslational modification, protein turnover, chaperones
C	216	4.49	Energy production and conversion
G	136	2.83	Carbohydrate transport and metabolism
E	299	6.22	Amino acid transport and metabolism
F	67	1.39	Nucleotide transport and metabolism
H	144	2.99	Coenzyme transport and metabolism
I	135	2.81	Lipid transport and metabolism
P	173	3.60	Inorganic ion transport and metabolism
Q	81	1.68	Secondary metabolites biosynthesis, transport and catabolism
R	350	7.28	General function prediction only
S	291	6.05	Function unknown
-	2,177	45.27	Not in COGs

^aThe total is based on the total number of protein coding genes in the annotated genome.

(LMG 28302) is a novel strain of *Bacillus azotoformans*, with similar redundancy in its nitrate reduction pathways, including the potential for DNRA, and a complete set of genes for oxic respiration and oxygen detoxification; its inability to grow with oxygen remains enigmatic.

Additional file

Additional file 1: Table S1. Overview of the genomic inventory for dissimilatory nitrogen transformations in *Bacillus azotoformans* MEV2011. **Table S2.** Overview of the genomic inventory for enzymatic reduction of O₂ and ATP synthase in *Bacillus azotoformans* MEV2011. **Table S3.** Overview of the genomic inventory for the detoxification of reactive oxygen species in *Bacillus azotoformans* MEV2011.

Abbreviation

DNRA: Dissimilatory nitrate reduction to ammonium

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

KF and AS designed research, MN isolated and characterized strain MEV2011 and carried out the genome sequencing, LS performed bioinformatics analyses, all authors analyzed data, MN and LS wrote the manuscript with help of AS and KF, all authors read and approved the final manuscript.

Acknowledgements

We thank Anne Stentebjerg and Britta Poulsen for their excellent technical assistance. Eline Palm Hansen is acknowledged for the isolation and initial characterization of strain MEV2011 (together with Maja Nielsen). This work was supported by the Technology Transfer Office (TTO), Aarhus University, Denmark. Funding for the Stellar Astrophysics Centre is provided by The Danish National Research Foundation (Grant agreement no.: DNRF106).

Author details

¹Section for Microbiology, Department of Bioscience, Aarhus University, Aarhus, Denmark. ²Center for Geomicrobiology, Department of Bioscience, Aarhus University, Aarhus, Denmark. ³Stellar Astrophysics Centre, Department of Physics and Astronomy, Aarhus University, Aarhus, Denmark.

Received: 24 June 2014 Accepted: 18 November 2014

Published: 21 January 2015

References

1. Fritze D. Taxonomy of the genus *Bacillus* and related genera: the aerobic endospore-forming bacteria. *Phytopathology*. 2004; **94**:1245–48.
2. Tiedje JM. Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In: Zehnder AJB, editor. *Biology of anaerobic microorganisms*. New York, NJ: John Wiley and Sons; 1988. p. 179–244.
3. Verbaendert I, Boon N, De Vos P, Heylen K. Denitrification is a common feature among members of the genus *Bacillus*. *System Appl Microbiol*. 2011; **34**:385–91.
4. Jones CM, Welsh A, Throbäck IN, Dörsch P, Bakken LR, Hallin S. Phenotypic and genotypic heterogeneity among closely related soil-borne N₂- and N₂O-producing *Bacillus* isolates harboring the *nosZ* gene. *FEMS Microbiol Ecol*. 2011; **76**:541–52.
5. Verbaendert I, De Vos P, Boon N, Heylen K. Denitrification in Gram-positive bacteria: an underexplored trait. *Biochem Soc Transac*. 2011; **39**:254–58.
6. Heylen K, Keltjens J. Redundancy and modularity in membrane-associated dissimilatory nitrate reduction in *Bacillus*. *Frontiers Microbiol*. 2012; **3**:371.
7. King E, Ward MK, Raney DE. Two simple media for the demonstration of pyocyanin and fluorescein. *J Laborat Clin Med*. 1955; **44**:301–07.
8. Pichinoty F, De Barjac H, Mandel M, Asselineau J. Description of *Bacillus azotoformans* sp. nov. *Int J Syst Bac*. 1983; **33**:660–62.
9. Spott O, Russow R, Stange CF. Formation of hybrid N₂O and hybrid N₂ due to codenitrification: First review of a barely considered process of microbially mediated N-nitrosation. *Soil Biol Biochem*. 2011; **43**:1995–2011.
10. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT. Gene Ontology: tool for the unification of biology. *Nat Genet*. 2000; **25**:25–9.
11. Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets. *Bioinformatics*. 2011; **27**:863–64.
12. Chevreux B, Wetter T, Suhai S. Genome sequence assembly using trace signals and additional sequence information. In: *Computer Science and Biology. Proceedings of the German Conference on Bioinformatics*. Hannover, Germany: GCB; 1999. p. 45–56.
13. Markowitz VM, Mavromatis K, Ivanova NN, Chen I-MA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics*. 2009; **25**:2271–78.
14. Bland C, Ramsey T, Sabree F, Lowe M, Brown K, Kyrpides N, Hugenholtz P. CRISPR Recognition Tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. *BMC Bioinformatics*. 2007; **8**:209.
15. Edgar R. PILER-CR: Fast and accurate identification of CRISPR repeats. *BMC Bioinformatics*. 2007; **8**:18.
16. Lowe TM, Eddy SR. tRNAscan-SE: A Program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res*. 1997; **25**:955–64.
17. Lagesen K, Hallin P, Rødland EA, Stærfeldt H-H, Rognes T, Ussery DW. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res*. 2007; **35**:3100–08.
18. Hyatt D, Chen G-L, LoCascio P, Land M, Larimer F, Hauser L. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinform*. 2010; **11**:119.
19. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res*. 1999; **27**:29–34.
20. Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S, Khanna A, Marshall M, Moxon S, Sonnhammer ELL, Eddy SR, Bateman A, Finn RD. The Pfam protein families database. *Nucleic Acids Res*. 2004; **32**(suppl 1):D138–41.
21. Tatusov R, Fedorova N, Jackson J, Jacobs A, Kiryutin B, Koonin E, Krylov D, Mazumder R, Mekhedov S, Nikolskaya A, Rao BS, Smirnov S, Sverdlov A, Vasudevan S, Wolf Y, Yin J, Natale D. The COG database: an updated version includes eukaryotes. *BMC Bioinformatics*. 2003; **4**:41.
22. Haft DH, Selengut JD, White O. The TIGRFAMs database of protein families. *Nucleic Acid Res*. 2003; **31**:371–73.
23. Auch AF, von Jan M, Klenk H-P, Göker M. Digital DNA-DNA hybridization for microbial species delineation by means of genome-to-genome sequence comparison. *Stand Genomic Sci*. 2010; **2**:117–34.
24. Bao P, Su J, Hu Z, Häggblom M, Zhu Y. Genome sequence of the facultative anaerobic bacterium *Bacillus* sp. strain ZYK, a selenite and nitrate reducer from paddy soil. *Stand Genomic Sci*. 2014; **9**:646–54.
25. Venkatraman R, Prakash O, Woyke T, Chain P, Goodwin LA, Watson D, Brooks S, Kostka JE, Green SJ. Genome sequences for three denitrifying bacterial strains isolated from a uranium- and nitrate-contaminated subsurface environment. *Genome Announc*. 2013; **1**(4):e00449–13. 10.1128/genomeA.00449-13.
26. Dilling W, Cypionka H. Aerobic respiration in sulfate-reducing bacteria. *FEMS Microbiol Letters*. 1990; **71**:123–27.
27. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, Ashburner M, Axelrod N, Baldauf S, Ballard S, Boore J, Cochrane G, Cole J, Dawyndt P, De Vos P, dePamphilis C, Edwards R, Faruque N, Feldman R, Gilbert J, Gilna P, Glöckner FO, Goldstein P, Guralnick R, Haft D, Hancock D. The minimum information about a genome sequence (MIGS) specification. *Nature Biotechnol*. 2008; **26**:541–47.
28. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A*. 1990; **87**:4576–79.
29. Skerman VBD, McGowan V, Sneath PHA. Approved Lists of Bacterial Names. *Int J Syst Bacteriol*. 1980; **30**:225–420.
30. Gibbons NE, Murray RGE. Proposals Concerning the Higher Taxa of Bacteria. *Int J Syst Bacteriol*. 1978; **28**:1–6.
31. Murray RGE. The Higher Taxa, or, a Place for Everything...? In: Holt JG, editor. *Bergey's Manual of Systematic Bacteriology*, Volume 1. 1st ed. Baltimore: The Williams and Wilkins Co.; 1984. p. 31–4.
32. List no. 132. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol*. 2010; **60**:469–72.
33. Ludwig W, Schleifer KH, Whitman WB. Class I. Bacilli class nov. In: De Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, Schleifer KH, Whitman WB, editors. *Bergey's Manual of Systematic Bacteriology*, Volume 3. 2nd ed. New York: Springer-Verlag; 2009. p. 19–20.
34. Prévot AR, Hauderoy P, Ehringer G, Guillot G, Magrou J, Prévot AR, Rosset D, Urbain A, editors. *Dictionnaire des Bactéries Pathogènes*. 2nd ed. Paris: Masson et Cie; 1953. p. 1–692.
35. Fischer A. Untersuchungen über Bakterien. *Jahrbücher für Wissenschaftliche Botanik*. 1895; **27**:1–163.
36. Cohn F. Untersuchungen über Bakterien. *Beitr Biol Pflanz*. 1872; **1**:127–224.
37. Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ, Kulam-Syed-Mohideen AS, McGarrell DM, Marsh T, Garrity GM, Tiedje JM. The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res*. 2009; **37**(suppl 1):D141–45.
38. Ludwig W, Strunk O, Westram R, Richter L, Meier H, Kumar Y, Buchner A, Lai T, Steppi S, Jobb G, Förster W, Brettske I, Gerber S, Ginhart AW, Gross O, Grumann S, Hermann S, Jost R, König A, Liss T, Lüßmann R, May M, Nonhoff B, Reichel B, Strehlow R, Stamatakis A, Stuckmann N, Vilbig A, Lenke M, Ludwig T. ARB: a software environment for sequence data. *Nucleic Acids Res*. 2004; **32**:1363–71.
39. Stamatakis A, Hoover P, Rougemont J. A rapid bootstrap algorithm for the RAxML web servers. *System Biol*. 2008; **57**:758–71.

doi:10.1186/1944-3277-9-23

Cite this article as: Nielsen *et al.*: Draft genome sequence of *Bacillus azotoformans* MEV2011, a (Co-) denitrifying strain unable to grow with oxygen. *Standards in Genomic Sciences* 2015 **9**:23.